

Infrared behaviour of the pressure in $g\phi^3$ theory in 6 dimensions

M. E. Carrington*

Department of Physics, Brandon University, Brandon, Manitoba, R7A 6A9 Canada

T. J. Hammond and R. Kobes†

Department of Physics, University of Winnipeg, Winnipeg, Manitoba, R3B 2E9 Canada

In an earlier paper Almeida and Frenkel considered the calculation of the pressure in $g\phi^3$ theory in 6 dimensions via the Schwinger–Dyson equation [1]. They found, under certain approximations, that a finite result ensues in the infrared limit. We find this conclusion to remain true with certain variations of these approximations, suggesting the finiteness of the result to be fairly robust.

I. INTRODUCTION

The infrared behaviour of gauge theories at finite temperature has long been a subject of interest. One aspect that has received particular attention is the fact noted by Linde that the thermodynamic pressure in Yang–Mills theory cannot be calculated perturbatively beyond 5th order in the coupling constant [2]. This problem has led to the development of a number of ways to extend the loop expansion in gauge theories at finite temperature, among which are approaches based on the Schwinger–Dyson equation [3, 4], the Braaten–Pisarski resummation scheme [5, 6], and a hybrid approach of perturbation and lattice gauge theory [7].

In an earlier paper Almeida and Frenkel used the Schwinger–Dyson equation to study the pressure in $g\phi^3$ theory in 6 dimensions [1]. This model is chosen because it has some qualitative similarities to Yang–Mills theory. They showed that when a certain set of ladder graphs are included the pressure is finite in the infrared limit. Given the fundamental importance of this conclusion, it is of some interest to examine to what degree the finite nature of the calculation is dependent on the approximations used. The purpose of this note is to study this question.

The paper is organized as follows. In Section II we set up a general framework for the calculation of the self–energy in the model, and show what approximations are used within this framework to obtain the results of Almeida and Frenkel [1]. Section III contains a discussion of four related variations of these approximations, which differ basically by including different approximations to the full propagator and vertex functions. We find that these different approximations lead to a quantitative difference with the results of Almeida and Frenkel [1], but the conclusion of the finite nature of the result in the infrared limit remains the same. Section IV contains some brief concluding remarks.

II. ALMEIDA–FRENKEL APPROXIMATIONS

The pressure in a field theory can be calculated by finding the self–energy function and using the fundamental relation:

$$\left(\frac{\delta P}{\delta \mathcal{D}_0} \right) = -\frac{T}{2} \Pi \quad (1)$$

where $\mathcal{D}_0$ is the free propagator and only one–particle–irreducible graphs are included in the self–energy function Π . To calculate this self–energy for ϕ^3 theory in 6 dimensions we consider the Schwinger–Dyson equation:

$$\Pi(p) = \frac{1}{2} \frac{g^2 T}{(2\pi)^5} \int_{\lambda}^T d^5 k \Gamma(p, k) G(k) G(p+k), \quad (2)$$

where λ is an infrared cutoff, $G^{-1}(p) = p^2 + \Pi(p)$ is the full inverse propagator, $\Gamma(p, k)$ is the full 3–point function, and we have taken the high temperature limit so that only the $n = 0$ term in the Matsubara frequency sum contributes.

*Electronic address: meg@theory.uwinnipeg.ca; Winnipeg Institute for Theoretical Physics, Winnipeg, Manitoba, R3B 2E9 Canada
 †Electronic address: randy@theory.uwinnipeg.ca; Winnipeg Institute for Theoretical Physics, Winnipeg, Manitoba, R3B 2E9 Canada

Consider now the approximation

$$\begin{aligned} G(p) &\approx \frac{1}{p^2} - \frac{\Pi(p)}{p^4} \\ \Gamma(p, k) &\approx 1 \end{aligned} \quad (3)$$

Under this assumption, Eq. (2) becomes

$$\Pi(p) = \frac{g^2 T}{48\pi^5 p^3} \left\{ \frac{1}{2} \int_{\lambda}^T k dk \log \left(\frac{k+p}{k-p} \right)^2 - \int_{\lambda}^T k dk \log \left(\frac{k+p}{k-p} \right)^2 \frac{\Pi(k)}{k^2} \right\} \quad (4)$$

If we now make the further approximations

$$\begin{aligned} \log \left(\frac{k+p}{k-p} \right)^2 &\approx \frac{4k}{p} & \dots k \ll p \\ &\approx \frac{4p}{k} & \dots k \gg p \end{aligned} \quad (5)$$

and define

$$\begin{aligned} \alpha &= \frac{g^2}{12\pi^3} \\ x &= \frac{\alpha T}{p} \\ y &= \frac{\alpha T}{k} \\ f(x) &= \frac{\Pi(p) - p^2}{p^2} \end{aligned} \quad (6)$$

Eq. (4) becomes

$$f(x) = -1 + \frac{x}{3} - \frac{x^2}{2\alpha} - x^2 \int_{\alpha}^x \frac{dy}{y^2} f(y) - x^4 \int_x^{\infty} \frac{dy}{y^4} f(y). \quad (7)$$

This equation was derived by Almeida and Frenkel [1], who show, analytically, that the solution for $f(x)$ is finite in the infrared limit $x \rightarrow \infty$. This result demonstrates that, using the approximations discussed above, the pressure is finite in the infrared limit.

III. OTHER APPROXIMATIONS

In this section we explore approximations other than Eqs. (3), (5) in solving the Schwinger Dyson equation (2). We consider two distinct questions: what happens if we don't make the approximation of Eq. (5) for the logarithmic terms, and what happens if we use a non-trivial 3-point interaction. As these approximations will require a numerical solution, in the following we choose $\alpha = 3$.

1. Approximation A

We first explore what happens if we don't make the approximations of Eq. (5) to the logarithmic term of Eq. (4). In this case, Eq. (4) can be written in the form

$$f(x) = -1 - \frac{x}{16\alpha^2} \left[(x^2 - \alpha^2) \log \left(\frac{x+\alpha}{x-\alpha} \right)^2 + 4\alpha x \right] - \frac{x^3}{4} \int_{\alpha}^{\infty} \frac{dy}{y^3} \log \left(\frac{x+y}{x-y} \right)^2 f(y) \quad (8)$$

in the infrared limit $\lambda \rightarrow 0$. Solving this equation numerically gives the same results as that found in [1] for the analytic solution of Eq. (7). These results appear in Fig. 1.




FIG. 1: Approximation A

The integral equation (8), and the equations that will be considered below, are subtle to solve, especially in the limit of large x (small momentum), which is the region we are mainly interested in. The source of the difficulty is as follows. In the limit $x \rightarrow \infty$ Eq. (8) has the form

$$f(x) = g(x) + x^3 \int_{\alpha}^{\infty} \frac{dy}{y^3} K(x, y) f(y), \quad (9)$$

This equation has a contribution from $g(x) \sim x^2$ which is canceled by a term coming from the integral. This cancellation is difficult to see numerically, but it can be avoided by performing the substitution,

$$f(x) = \tilde{f}(x) + \frac{\alpha^2 \sigma}{x^2} + f_{\infty} \quad (10)$$

with constants σ and f_{∞} to be fixed shortly. Using this substitution, we find that $\tilde{f}(x)$ satisfies an equation of the form

$$\tilde{f}(x) = \tilde{g}(x, \sigma, f_{\infty}) - \frac{\alpha^2 \sigma}{x^2} - f_{\infty} + x^3 \int_{\alpha}^{\infty} \frac{dy}{y^3} K(x, y) \tilde{f}(y) \quad (11)$$

We fix σ by requiring $\tilde{g}$ approach a constant as $x \rightarrow \infty$, and we fix f_{∞} by requiring the solution $\tilde{f}(x)$ approach 0 as $x \rightarrow \infty$.

2. Approximation B

In this section and the next we use an eikonal approximation for the 3-point function [8, 9, 10]:

$$\Gamma(p, k) \approx 1 + \frac{\Pi(k+p) - \Pi(k)}{\vec{p}^2 + 2\vec{k} \cdot \vec{p}} \quad (12)$$

Using this expression effectively resums a series of higher loop vertex graphs which are important in the infrared region. First, we consider the integral equation obtained from Eq. (2) with the vertex given by Eq.(12) and bare propagators:

$$G(p) \approx \frac{1}{p^2} \quad (13)$$

Using the notation of Eq. (6) and taking the limit $\lambda \rightarrow 0$ we obtain,

$$f(x) = -1 + \frac{x}{4\alpha^2} \left[\frac{3}{4}(x^2 - \alpha^2) \log \left(\frac{x + \alpha}{x - \alpha} \right)^2 - \frac{1}{8}(4x^2 - \alpha^2) \log \left| \frac{2x + \alpha}{2x - \alpha} \right| + \frac{5}{2}\alpha x \right] \\ - \frac{x^3}{4} \int_{\alpha}^{\infty} \frac{dy}{y^3} \left\{ \frac{3}{2} \log \left(\frac{x + y}{x - y} \right)^2 - \log \left| \frac{2x + y}{2x - y} \right| \right\} f(y) \quad (14)$$

The results, which appear in Fig. 2, indicate that the self-energy function is finite in the infrared limit $x \rightarrow \infty$, although, not surprisingly, we obtain an asymptotic value quantitatively different from that of Fig. 1.




FIG. 2: Approximation B

3. Approximation C

We next consider the integral equation obtained from Eq. (2) using

$$G(p) \approx \frac{1}{p^2} - \frac{\Pi(p)}{p^4} \\ \Gamma(p, k) \approx 1 + \frac{\Pi(k + p) - \Pi(k)}{\vec{p}^2 + 2\vec{k} \cdot \vec{p}} \quad (15)$$

Using the notation of Eq. (6) and taking the limit $\lambda \rightarrow 0$ we obtain,

$$f(x) = -1 + \frac{x}{4\alpha^2} \left[\frac{1}{4}(x^2 - \alpha^2) \log \left(\frac{x + \alpha}{x - \alpha} \right)^2 - \frac{1}{8}(4x^2 - \alpha^2) \log \left| \frac{2x + \alpha}{2x - \alpha} \right| + \frac{1}{2}\alpha x \right] \\ - \frac{x^3}{4} \int_{\alpha}^{\infty} \frac{dy}{y^3} \log \left| \frac{2x + y}{2x - y} \right| f(y) \quad (16)$$

These results appear in Fig. 3. As before, a finite result in the infrared limit is obtained. The fluctuations that occur at small x are not physical since small x corresponds to the ultraviolet limit, and the eikonal approximation of Eq. (12) is only valid in the infrared limit.




FIG. 3: Approximation C

4. Approximation D

The final approximation we consider is a variation of the previous one. We use the eikonal expression Eq. (12) for the vertex and work with full propagators of the form: $G(p) = 1/p^2 + \Pi(p)$. The Schwinger–Dyson equation becomes:

$$\Pi(p) = \frac{g^2 T}{(2\pi)^5} \int_{\lambda}^T d^5 k \frac{G(k)}{\vec{p}^2 + 2\vec{k} \cdot \vec{p}} \quad (17)$$

At this point we make the approximation

$$G(p) \approx \frac{1}{p^2} - \frac{\Pi(p)}{p^4} \quad (18)$$

and obtain, in the limit $\lambda \rightarrow 0$,

$$f(x) = -1 - \frac{x^3}{4} \int_{\alpha}^{\infty} \frac{dy}{y^3} \log \left| \frac{2x+y}{2x-y} \right| f(y) \quad (19)$$

The difference between Eq. (19) and Eq. (16), which was obtained using Approximation C, is as follows. Eq. (19) makes use of an analytic cancellation between certain sets of graphs corresponding to vertex corrections, which are contained in Eq. (12), and graphs that are contained in the expression for the full propagator $G(p)$ [10]. These analytic cancellations are performed, before the expansion of the full propagator is done. The results appear in Fig. 4. Once again we obtain a self-energy that is finite in the infrared limit.




FIG. 4: Approximation D

IV. CONCLUSIONS

We have considered various approximations to the Schwinger–Dyson equation that can be used to calculate the pressure in $g\phi^3$ theory in 6 dimensions. The approximations used sum up different classes of diagrams believed to be important in the infrared limit. In all cases considered a finite result is obtained for the pressure in the infrared limit, although the quantitative behaviour differs according to which approximation is used, as is expected. The important conclusion to be drawn from these results is that the finiteness of the pressure in the infrared limit is not very sensitive to the particular approximation used, and thus seems to be a general feature of pressure calculations based on a Schwinger–Dyson approach.

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] A. P. de Almeida and J. Frenkel, Phys. Rev. **D47**, 640 (1993).
 - [2] A. D. Linde, Phys. Lett. **96B**, 289 (1980).
 - [3] O. K. Kalashnikov and V. V. Klimov, Fortschr. Phys. **31**, 613 (1983).
 - [4] K. Kajantie and J. Kapusta, Ann. Phys. (N. Y.) **160**, 477 (1985).
 - [5] E. Braaten and R. Pisarski, Nucl. Phys. **B337**, 569 (1990).
 - [6] J. Frenkel, A. V. Saa, and J. C. Taylor, Phys. Rev. **D46**, 3670 (1992).
 - [7] E. Braaten and A. Nieto, Phys. Rev. **D53**, 3421 (1996).
 - [8] J. M. Cornwall and G. Tiktopoulos, Phys. Rev. **D15**, 2937 (1977).
 - [9] J. M. Cornwall and W.-S. Hou, Phys. Lett. **B153**, 173 (1985).
 - [10] M. Carrington, R. Kobes and E. Petitgirard, Phys. Rev. **D57**, 2631–2634 (1998).